

Upper limit on the $\eta \rightarrow \pi^0 \pi^0$ decay

M.N. Achasov, A.V. Berdyugin, A.V. Bozhenok, D.A. Bukin,
S.V. Burdin, T.V. Dimova, S.I. Dolinskii, V.P. Druzhinin,
M.S. Dubrovin, I.A. Gaponenko, V.B. Golubev, V.N. Ivanchenko,
A.A. Korol, S.V. Koshuba, I.N. Nesterenko, E.V. Pakhtusova,
E.A. Perevedentsev, A.A. Salnikov, S.I. Serednyakov, V.V. Shary,
Yu.M. Shatunov, V.A. Sidorov, Z.K. Silagadze*, Yu.S. Velikzhanin.

Budker Institute of Nuclear Physics
630 090, Novosibirsk, Russia

Abstract

Upper bound $\text{Br}(\eta \rightarrow \pi^0 \pi^0) < 6 \cdot 10^{-4}$ with 90 % confidence is reported from the SND experiment at the VEPP-2M collider.

CP-noninvariance still remains the most mysterious phenomenon in elementary particle physics because it implies either an absolute difference between left and right in nature, or maybe even more exotic possibility of the mirror world existence [1].

CP-violation is incorporated into the Standard Model by means of complex coupling constants in the quark mixing matrix and is controlled by a single parameter - the Kobayashi-Maskawa phase [2]. The only manifestations of the CP violating interactions, observed by now, are restricted to the K_L meson decays. More profound CP-violating effects are predicted in the B^0 -meson decays but the corresponding experiments are still awaiting their time.

Flavor-conserving CP-violating interactions, like $\eta \rightarrow 2\pi$ decay, are expected to be extremely small in the Standard Model, namely $\text{Br}(\eta \rightarrow \pi^+ \pi^-) \leq 2 \cdot 10^{-27}$ [3, 4]. Nevertheless, experimental search for the $\eta \rightarrow 2\pi$ decay is very important [5], because unconventional CP-violating effects have "a golden opportunity" to reveal themselves in this decay.

Let us mention two of such CP-violation models beyond the Standard Model. $\eta \rightarrow 2\pi$ decay can take place if there exists a strong CP-violation

*Corresponding author. Fax +7 3832 35 21 63, e-mail silagadze@inp.nsk.su

due to QCD Θ -term [6]. Using present experimental constraint on this Θ -term from the electric dipole moment of the neutron, one can obtain $\text{Br}(\eta \rightarrow \pi^+\pi^-) \leq 0.78 \cdot 10^{-16}$ [3, 4]. Another possibility is a spontaneous CP violation in the extended Higgs sector [7]. It was shown in [4] that in this case $\text{Br}(\eta \rightarrow \pi^+\pi^-)$ can be as high as $1.2 \cdot 10^{-15}$. But even this last value is too small for $\eta \rightarrow 2\pi$ decay to be observed in any present or planned experiments. So its observation will be a clear signal of some CP-violation exotica.

The only experimental upper limit $\text{Br}(\eta \rightarrow \pi^+\pi^-) \leq 1.5 \cdot 10^{-3}$ was reported in [8]. Recently this result was confirmed by CMD-2 detector [9]. Here we present, for the first time, upper bound $\text{Br}(\eta \rightarrow \pi^0\pi^0) < 6 \cdot 10^{-4}$ obtained in SND experiment at VEPP-2M e^+e^- collider.

SND [10] is a general purpose nonmagnetic detector. Its main part, very important for neutral decay channels, is a highly segmented three layer NaI(Tl) calorimeter [11] of 3.6 t total weight with vacuum phototriode [12] readout.

During 1996 experimental run, SND collected data corresponding to integrated luminosity of about 3.9 pb^{-1} . The data sample consists of six successive scans of the narrow energy interval covering the ϕ -meson peak (FI96 experiment [13]). Total number of produced ϕ -mesons is $7.4 \cdot 10^6$, corresponding to $9.3 \cdot 10^4$ η -mesons, produced through $\phi \rightarrow \eta\gamma$ decay. Five-photon events from this data sample, with five clusters in the calorimeter and no charged particles, were used to search for a signal from the $\eta \rightarrow 2\pi^0$ decay, in the reaction $\phi \rightarrow \eta\gamma, \eta \rightarrow 2\pi^0$. Only the events having the normalized full momentum less than 0.15, and the normalized total energy deposition in the calorimeter greater than 0.8 were selected for the analysis.

For events which had two π^0 candidates with

$$(m_{2\gamma}^{(1)} - m_{\pi^0})^2 + (m_{2\gamma}^{(2)} - m_{\pi^0})^2 < 300\text{MeV}^2$$

a kinematical fit was performed assuming energy-momentum conservation and presence of two π^0 mesons in the final state. χ^2 of this kinematical fit can be used to reject background from the $\phi \rightarrow K_S K_L$ decay, because these background events have a poor energy-momentum balance. In the present analysis we used mild cut $\chi^2 < 40$.

Another background comes from the $\phi \rightarrow \eta\gamma, \eta \rightarrow 3\pi^0$ decay. Examining the transverse energy profile of the electromagnetic shower in the calorimeter, one can recognize and reject events where close photons

merge together [14]. This merging is one of the main reasons why $3\pi^0\gamma$ final state can mimic $2\pi^0\gamma$ final state. We demand the corresponding separation parameter (described in detail in [14]) to be less than 5, which is also a mild cut.

Remaining events, surviving these cuts, are mainly due to $e^+e^- \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$ reaction. To suppress this background, additional cuts $m_{\pi^0\gamma} < 730\text{MeV}$ and $\chi_\omega^2 > 20$ were applied. Here χ_ω^2 is a parameter describing degree of likelihood of the assumption that our 5-photon event is a $e^+e^- \rightarrow \omega\pi^0 \rightarrow \pi^0\pi^0\gamma$ event.

The photon recoil mass distribution for remaining 70 events is shown in fig.1. Note, that Monte-Carlo simulation predicts that 11 ± 3 background events from the $e^+e^- \rightarrow \omega\pi^0$ and 19 ± 5 events from the $e^+e^- \rightarrow \eta\gamma \rightarrow 3\pi^0\gamma$, 3γ , ... reactions surviving these cuts. It is difficult to estimate the remaining background from the $\phi \rightarrow K_S K_L$ channel because of huge Monte-Carlo statistics and high accuracy for the K -mesons nuclear interactions simulation needed for this goal. No candidate events passed analysis cuts from the simulated $4 \cdot 10^5$ $\phi \rightarrow K_S K_L$ decays.

It turns out that the most significant background comes from the $\phi \rightarrow f^0\gamma$, $f^0 \rightarrow 2\pi^0$ channel: 52 ± 13 events are expected according to MC, if the recently published parameters of this decay [15] are used in simulation.

Background from the $\phi \rightarrow \eta\gamma$, $\eta \rightarrow \pi^0\gamma\gamma$ decay seems to be negligible. Only 0.7 ± 0.2 events are expected due to very small $\eta \rightarrow \pi^0\gamma\gamma$ branching ratio.

In total we expect 83 ± 14 background events from the processes described above, which is in a good agreement with the experimentally observed number. Although, in the $\pm 3\sigma$ interval around the expected η -peak only one experimental event was observed, while Monte-Carlo predicts 10 ± 5 events. Due to small efficiencies for background processes in the signal area, significant systematic errors could be expected in their simulation. So we don't perform any background subtraction and our result is based only on the experimental statistics.

Fig.1 shows that no significant signal from the $\eta \rightarrow 2\pi^0$ could be seen over the expected background from the $f^0\gamma$, $\omega\pi^0$ and $\eta \rightarrow 3\pi^0$ channels. Taking $\epsilon = 7.5\%$ as a detection efficiency for the $\eta \rightarrow 2\pi^0$ decay, which follows from the Monte-Carlo analysis, and assuming $N_x = 3.89$ for Poisson upper limit, which corresponds to the 90 % confidence level

for one observed event [16], we get the upper bound

$$\text{Br}(\eta \rightarrow 2\pi^0) < \frac{N_x}{N_\eta \cdot \epsilon} \approx 6 \cdot 10^{-4} \ .$$

The work is partially supported by RFBR (Grants No 96-02-19122, 97-02-18563).

References

- [1] T. D. Lee and C. N. Yang, Phys. Rev. **104**, 254 (1956).
I. Yu. Kobzarev, L. B. Okun, and I. Ya. Pomeranchuk, Yad. Fiz. **3**, 1154 (1966).
R. Foot, H. Lew and R. R. Volkas, Phys. Lett. **B272**, 67 (1991).
R. Foot and R. R. Volkas, Phys. Rev. **D52**, 6595 (1995) .
Z. K. Silagadze, Phys. At. Nucl. **60**, 272 (1997) .
- [2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973) .
- [3] C. Jarlskog and E. Shabalin, Phys. Rev. **D52**, 248 (1995) .
- [4] C. Jarlskog and E. Shabalin, Phys. Rev. **D52**, 6327 (1995) .
E. Shabalin, in “*Second DAΦNE Physics Handbook*”, edited by L. Maiani, L. Pancheri and N. Paver (INFN, Frascati, 1995) .
- [5] I. Yu. Kobzarev and L. B. Okun, Zh. Eksp. Teor. Fiz. **46**, 1418 (1964) .
- [6] R. J. Crewther, P. Di Vecchia, G. Veneziano, E. Witten, Phys. Lett. **88B**, 123 (1979) .
M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B166**, 494 (1980) .
- [7] S. Weinberg, Phys. Rev. Lett. **37**, 657 (1976) .
T. D. Lee, Phys. Rev. **D8**, 1226 (1973) .
- [8] J. J. Thaler, J. A. Appel, A. Kotlewski et al., Phys. Rev. **D7**, 2569 (1973) .
- [9] R. R. Akhmetshin et al., Proc. of the 28-th International Conference on High Energy Physics, Warsaw, Poland, 25-31 July, 1996. p. 522.

- [10] V. M. Aulchenko et al., Budker INP 95-56, Novosibirsk, 1995.
V. M. Aulchenko et al., Proc. Workshop on Physics and Detectors for DAΦNE, Frascati, April 1991, p. 605.
- [11] A. D. Bukin, S. I. Dolinsky, V. P. Druzhinin et al.,
Phys. At. Nucl.**56**, 1494 (1993) (Yad. Fiz. **56**, 75 (1993)) .
M. N. Achasov, A. D. Bukin, D. A. Bukin et al., Nucl. Instr. and Meth. **A 379**, 505 (1996) .
- [12] P. M. Beschastnov, V. B. Golubev, E. A. Pyata et al., Nucl. Instr. and Meth. **A 342**, 477 (1994) .
V. V. Anashin, P. M. Beschastnov, I. I. Goldberg et al., Nucl. Instr. and Meth. **A 379**, 522 (1996) .
- [13] M.N.Achasov, M. G. Beck, P. M. Beschastnov et al., Novosibirsk preprint BUDKERINP-96-47 .
- [14] M. G. Bekishev, V. N. Ivanchenko, Nucl. Instr. and Meth. **A 361**, 138 (1995) .
A. V. Bozhenok, V. N. Ivanchenko, Z. K. Silagadze, Nucl. Instr. and Meth. **A 379**, 507 (1996) .
- [15] M.N. Achasov, M.G. Bek, K.I. Beloborodov et al., Novosibirsk preprint BUDKERINP-97-78 (hep-ex/9710017).
M.N. Achasov, A.V. Berdyugin, A.V. Bozhenok et al., Evidence of $\phi \rightarrow \pi^0 \pi^0 \gamma$ and $\phi \rightarrow \pi^0 \eta \gamma$ decays in SND experiment at VEPP-2M. Talk given at HADRON-97 Conference, BNL, August 25-30, 1997 (hep-exp/9711023).
- [16] Particle Data Group, Phys. Rev. **D54**, 164 (1996) . p. 164.




Figure 1: Photon recoil mass. a) - experiment. Solid curve indicates the position and the width (according to MC simulation) of a possible signal from the $\eta \rightarrow 2\pi^0$ decay. The peak amplitude corresponds to the $3 \cdot 10^{-3}$ branching ratio. b) - expected background according to MC simulation.